arXiv:astro-ph/0503012v1 1 Mar 2005

Neutrino Absorption: In The Magnetic Field Of GRB In The Fireball Model.

A. K. Ganguly^{a*}
Wintel Communications, Kolkata 700030
India.

November 2004

Abstract

TeV or Super TeV neutrinos are expected to originate at the Gamma ray burster (GRB) events in the universe. These neutrinos are expected to be produced from the photo-meson interaction of the protons in the GRB environment. In the usual picture, the protons in the GRB fireball undergo Fermi acceleration in the ambient magnetic field of the GRB to very high energy. These protons then interact with the MeV, KeV or meV photons to produce Delta particles those subsequently decay to produce very high energy neutrinos. In this note we focus on the 100 TeV neutrinos produced in a GRB in the fireball model. We try to estimate the production point of these multi TeV neutrinos from the center. The strength of the Magnetic field there. Lastly but importantly modification in the spectrum because of neutrino absorption in the ambient magnetic field. The strength of the ambient magnetic field may change the neutrino spectrum.

1 Introduction

It is established now that GRB's are among the most violent energy releasing events of the universe. Total energy release in these events could be equivalent to few solar mass and are seen to be distributed isotropically all around the universe. They are very high redshift objects occurring in nature – implying their cosmological origin. Since their discovery, the physics of the process that powers these events is yet to be understood. For some time now it has been conjectured that GRBs are source of very high energy cosmic rays. Moreover they have also been tipped off as one of the source of very high energy neutrino. The energy reachable in the GRB mediated events are several orders of magnitude higher than those available in the laboratory, that is one of the motivations why there is some interest in the community to study the physics of GRB.

The origin of high energy particles in a GRB is ascribed to Fermi mechanism in the ambient magnetic field in the GRB enviornment. As we discuss later in this paper the same magnetic field might be responsible for modifying the ν spectra. This requires good understanding of GRB physics as well as, careful evaluation of GRB parameters like temperature, Magnetic field, jet Lorentz Boost factor, matter density

---

[*] aviavi85@rediffmail.com

in the rest frame of the jet etc. We discuss them below one after the other.

As already mentioned, GRB's are very high redshift objects, implying, they must have happened during the time of formation of the first generation stars of our universe. They have certain distinct features, such as, the rapid variability time (10^{-2} Sec), the non-thermal nature of the spectra, the hardness of the spectra ($\sim$ 100 MeV). The existence of large energy in the power law tails of the GRB spectra— above threshold gives rise to whats usually referred in the literature as *compactness problem* [1]. The naive estimate of the source size from the variability of the spectra runs in to problem. This is principally because of the fact, that a stationary compact object with an energy release, of the order of 10^{52} erg/sec would produce an optically thick environment; however the fact that one observes non-thermal spectra, gave rise to what is called *Compactness Problem Of GRB*. The time variability and hardness of the spectra together suggest the source to be extremely compact in size ($r_o \sim 10^7$ cm), from which a highly energetic jet moves outward around the rotational axis of the object.Actually the unusually hard and non-thermal nature of the photon spectra suggest that the jet must be moving towards the observer with high Lorentz boost Γ, whose magnitude can vary between several hundreds to few tens of thousands.

Inside the moving jet, the medium has appropriate optical thickness to last the radiation in the jet up to the variability time (of the jet). That could vary between 1 to 10^{-4} sec. To have the jet optically thick to (One to two MeV photons), it has to have some baryonic contamination in – composed of p and n, [2]. They are accelerated to very high energy in the shock and ambient magnetic field of the GRB through Fermi mechanism and subsequently would produce pions through photo-meson or internal collision processes. These pions finally decay to high energy neutrinos, electrons and positrons. In this note we would be focusing mainly on neutrinos.

The neutrinos produced in GRB, are emitted at various stages of the jet evolution with varying energy. Starting from low energy neutrinos at the beginning to extremely high energy neutrino energy at the end. Typically GeV neutrinos are expected to be produced at the initial stage, followed by 5 TeV or more and then $\geq$ 100 TeV ones.

Neutrino properties are known to be sensitive to ambient Electro-magnetic field and matter field [3]. In particular there are several decay channels, forbidden in vacuum but are allowed in a background magnetic field. Actually these are the motivating reasons to look into aspects of the magnetic field structure, medium properties; at locations where neutrinos of specific energy would originate. With this purpose in mind, in this note we would estimate the radial distance form the center of the star (r_π) where the energetic pions form to decay into 100 TeV neutrinos. We then go ahead to estimate the magnetic field there, followed by estimates of magnetic field induced neutrino decay.

1.1 Neutrino Production In GRB.

As has been mentioned in the beginning, in this note we investigate ultra high energy neutrino production and their interactions in the ambient medium and the possibility of change in the spectrum because of that. The reaction dynamics is as follows, pions first decay to muon and muon type neutrino, i.e.,

$$\pi^+ \rightarrow \mu^+ + \nu_\mu, \tag{1.1}$$

followed by the muons decaying to,

$$\mu^+ \rightarrow e^+ + \nu_e + \bar{\nu}_\mu. \tag{1.2}$$

From Eq.(1.1) and (1.2) its clear that, $\nu_\mu \& \bar{\nu}_\mu$ emission takes place from two different reactions. The first emission would be taking place within a radius (r_π, the pionosphere radius) within which the pions produced from photo meson interaction would decay. The other region being the location where the energetic muons decay (usually referred in literature as r_d). For a given r_d one can derive the energy of the muon anti-neutrinos observed on earth ($\epsilon^{ob}_{\bar{\nu}\mu}$) or vice versa. The muon anti-neutrino in Eq. (1.2) is an end product of a three body decay, therefore the energy carried by the $\bar{\nu}_\mu$, in the rest frame of the muon– would be $\frac{1}{3}m_\mu$ (roughly!). Where m_μ is the rest mass of the muon. Hence the anti-neutrino energy as observed on earth would be given by,

$$\epsilon^{ob}_{\bar{\nu}\mu} = \frac{1}{3}m_\mu \Gamma_\mu \Gamma_h, \quad \text{In units of } \hbar\text{=c=1}. \tag{1.3}$$

Where Γ_h is the Lorentz boost of the wind with respect to an observer on earth. On the other hand, if the muons with half life t_μ originate in the jet wind of the GRB (extending up to distance r_d), then it follows from there that,

$$r_d = \Gamma_\mu \Gamma_h t_\mu. \tag{1.4}$$

From relations Eq.(1.3) and (1.4) one can relate the observed anti-neutrino energy with that of the wind dynamical radius, i.e.,

$$\epsilon^{ob}_{\bar{\nu}\mu} = \frac{m_\mu r_d}{3t_\mu}. \tag{1.5}$$

For the muon type neutrinos, produced in a frame where the pion is at rest, the available kinetic energy is roughly close to $(m_\pi - m_\mu) = 34$ MeV and the resulting muon is non-relativistic. So even if most of the energy is carried by the ν_μ it would not be very different from the $\bar{\nu_\mu}$ or the ν_e, i.e $\sim \frac{1}{3}m_\mu$. Although their point of origin in the GRB environment would vary (because of difference in lifetime of pion or muon), hence change in the Lorentz Boost, however given the detector efficiency, that probably would not be detectable. Therefore we would consider the point of their origin to be same.

We would like to note, in absence of magnetic field the ratio of number density of neutrinos $(n_{\nu_e} n_{\nu\mu})$ would be 1:1, at the production point. However in a magnetic background new decay channels, e.g., $\nu \to \nu l^+ l^-$ opens up with nonzero probability α. In general α depends of lepton mass m_l their energy E_ν, strength of Magnetic field B etc. Since these many of these quantities are flavor dependent, implying different absorption rate for different neutrino flavors. And to find out the absorption rates we need to estimate the parameters like Strength of Magnetic Field at a particular location. The energy of the neutrinos being produced there etc. In view of this this has been estimated in this paper, following the fireball model of [2].

To estimate various physical parameters of a GRB, we would stick to the fireball model of GRB (successful) and try to refrain from making any explicit assumption about the progenitor. The postulates and assumptions we make, would be based mostly on [2], [7], [5] and [6].

The organization of this document is as follows, in section II, following fireball model we discuss the physics of GRB. Section III is devoted to estimating various parameters, and lastly section IV would deal with details of absorption rate and the asymmetry in number density of ν_e and ν_μ for some specific energy ≥ 100 TeV. we conclude by discussion and some numerical estimates of asymmetry.

2 Physics Of GRB

Numerical simulations based studies reveal that, the jet produced at the central engine of the GRB moves along the rotation axis with a Lorentz factor (Lf) Γ. After launch.jet consists of shocked jet material and shocked stellar materials. Initially, the jet heats up the region where it interacts with the stellar material, till the pressure in this zone becomes comparable to the same in the jet. Following notations of [4] region ahead of shock frame is called up stream, the one behind the shock called down stream. As the pressure in the up stream termination shock grows material flows into the jet creating a reverse shock (Lorentz factor Γ_R). Subsequently material in the shocked region down stream (jet head) move out ward with Lorentz factor Γ_h. The synchrotron and inverse Compton is believed to diffuse in, thermals and power the movement of the jet head. The isotropic luminosity of the GRB usually is related with the temperature of the medium in the jet head.In the rest frame of the jet (of Lf, Γ_h) the retarded jet moves with a Lorentz factor $\frac{\Gamma_j}{2\Gamma_h}$ [7].

After injection, during its passage through C/O/He core jet head moves with subreltivistic velocity. However beyond He core there is a significant drop in the density of the stellar material (envelope)i.e., $\rho \leq 10^{-7}$ gm/cc. and the jet can accelerate. As will be discussed below, during this phase $\Gamma_h \propto \frac{r}{r_o}$, where r and r_o are the radial distance at the point of concern and the injection point. The jet accelerates till it's L.f reaches a saturation value. Further out it follows a self-similar motion and decelerates while interacting with the (ISM);we note here that baryonic components (mainly protons) of the jet are accelerated to high energy in the reverse shock and these high energy protons interacting with the thermal photons produce neutrinos.

Near the outer envelope, jet is believed to be dissipation-less hence the jet fluid in GRB can be collisionless. It behaves like an ideal Magneto Hydro Dynamic (MHD) fluid. As a result it follows that the entropy and the energy of the fluid component can be held constant. Conservation of entropy and energy implies [9],

$$r^2\Gamma(r)r_oT_\gamma^3(r) = \text{constant.} \tag{2.1}$$

and

$$r^2\Gamma(r)r_o\Gamma(r)T_\gamma^4(r) = \text{constant.} \tag{2.2}$$

In the equations above, r_o stands for radius at the injection point and its usually taken as $r_o = 10^7 cm$. We would like to point out that, $\Gamma(r)$ the position dependent Lorentz factor, from now on, would be replaced by Γ_h unless explicitly mentioned. From the estimates of Luminosity in terms of temperature one obtains, the estimate of the temperature at the injection point turns out to be,

$$T_o = 1.2L_{52}^{1/4}r_{o7}^{-1/2}\text{Mev.} \tag{2.3}$$

With $L_{52} = 10^{-52}L_{iso}$ erg/sec and $r_o = 10^{-7}r_{07}$ cm. Assuming, $\Gamma(r)|_{r=r_o} = O(1)$, it is possible to estimate the rhs of eqns. (2.1) and (2.2). They turn out to be,

$$r^2\Gamma(r)r_oT_\gamma^3(r) \quad \sim \quad 2.1\times10^{53}L_{52}^{3/4}. \tag{2.4}$$

and

$$r^2\Gamma(r)r_o\Gamma(r)T_\gamma^4(r) \quad \sim \quad 2.5\times10^{53}L_{52}\text{MeV.} \tag{2.5}$$

As before unless explicitly mentioned we would suppress the explicit position dependence of temperature $T(r)$. Relations (2.4), (2.5) imply,

$$\Gamma_h \propto r. \tag{2.6}$$

Assuming the presence of only magnetic field, ((eB), in the rest frame of the plasma.), using dissipationless fluid equations, it is possible to derive an exact relation,

$$\left(\frac{eB}{2}\right)^2 = n_jm_p. \tag{2.7}$$

Where, we assumed the medium to be composed of heaviest charged particle available in the plasma, namely protons . Their number per unit volume in jet rest frame is defined as n_j. This is justified because the contribution from electrons would be far too less. Further more if we assume them to carry the isotropic energy (luminosity), L ergs/sec, then the number density of protons, at some distance r (to an observer on earth) from the center of the GRB (moving with Lf, Γ_j) towards earth would be [6],

$$n_j = \frac{\mathrm{L}}{4\pi r^2 \Gamma_j^2 m_p}. \tag{2.8}$$

For a locally charge neutral plasma, Eq.(2.8) would also provide the number density of electrons.

The physics of the shock front is described by Hugoniot equations.They balance various thermodynamic quantities like, energy, pressure, etc., across the discontinuity of the shock front. For instance equating the pressure between the forward and backward shock, (with density of material ρ in units of gm cm^{-3}) it has been shown [7] that ,

$$\begin{aligned} \Gamma_h^4 &= \frac{\Gamma_j^2}{4\rho} n_j m_p \\ &= \frac{L}{16\pi r^2 \rho}. \end{aligned} \tag{2.9}$$

Following [8] near the outer envelope one can parametrize $\rho = \rho_* \left(\frac{R}{r} - 1\right)^3$ with $\rho_* \leq 10^-7$gm/cc. Using this parametrization, it turns out that,

$$\Gamma_h = \frac{Lr}{16\pi\rho_* \left(R - r\right)^3} \tag{2.10}$$

which matches with the suggested form $\Gamma_h \propto r$, if one expands the denominator in powers or $\frac{r}{R}$, for $r < R$ and retains the leading order term.

3 Parameters

3.1 Temperature Of the plasma

In this section we would try to estimate the temperature, that the shocked plasma jet moves with. Since photons are believed [2] to be radiated from the shock heated plasma jet, moving with Lf Γ_h then all the radiated photons from the GRB would account for its total luminosity $\mathrm{L_{iso}}$. Using the emissivity-temperature relationship for pure photon gas it is possible to establish the temperature of the emitted radiation. The same under this assumption turns out to be,

$$T_r \sim 10 \left(\frac{\mathrm{L_{iso}}}{r^2 \Gamma_h^2}\right)^{1/4}, \tag{3.1}$$

However this relation is bound to get multiplied by correction factors (< 1) if the photons constitute a part of the total luminosity L. However to keep the physics simple we assume the photons to carry the total energy of the GRB. We conclude this section by noting that, the number density of the photons at this temperature would be given by.

$$n_\gamma = \mathrm{T_r^3}. \tag{3.2}$$

which essentially follows the laws of pure photon gas. One can use eqn. (3.1) to express photon number density in terms of total luminosity L and radial depth from the center.

3.2 r_π: Pion Formation Radius

In previous sections we have estimated the various parameters related to the production of high energy protons as well as the approximate temperature profile of the shocked plasma head. In view of these estimates it is appropriate to estimate the the distance from the origin where these protons (of energy ϵ_p) would scatter off the thermal photons to undergo Δ formation. In the jet head, the protons at the resonant energy would scatter off thermal photons to produce Δ as long as,at the rest frame of the jet, the following energy condition, in obvious notation, is given by,

$$\epsilon_\gamma \epsilon_p = 0.2\mathrm{GeV}^2. \tag{3.3}$$

Before proceeding further we note that, the reaction rate R for $p + \gamma \to \Delta$ would be given by

$$R = n_\gamma n_j \sigma_{p,\gamma\to\Delta} \left[1 - cos\left(\theta\right)\right] \sim n_\gamma n_j \sigma_{p,\gamma\to\Delta}, \tag{3.4}$$

Here we have assumed $\theta = 90^o$. Its appropriate to note that, in order to have resonant photon proton scattering to Δ, inside a GRB shock cavity, number of protons (n_j) in unit volume should be equal to $R \times \delta t$; with δt being variability time for the GRB, this leads to the condition,

$$n_\gamma \sigma_{p\gamma\to\Delta} \delta t = 1. \tag{3.5}$$

In eqn.(3.5) above, $\sigma_{p\gamma\to\Delta} = 5\times10^{-28}\mathrm{cm}^2$, is the resonant proton photon cross section for Δ production. Eqn. (3.2) implies at a distance r from the center, $n_\gamma = \frac{L_{iso}}{4\pi r^2 \Gamma_h^2} T_\gamma$, where $T_\gamma = \epsilon_\gamma$, i.e., the energy of the photon. Using this relation, eqn.(3.5) implies,

$$\frac{L_{iso}\ \sigma_{p\gamma\to\Delta}\ \Delta t}{4\pi\ r_h^2 \Gamma_h^2\ \epsilon_\gamma} = 1. \tag{3.6}$$

In order to have resonance scattering, the energy condition, i.e., eqn. (3.3) should be satisfied. Replacing ϵ_γ in terms of ϵ_p by using, (eqn. (3.3)) and the expression for Γ_h from eqn. (2.9), one arrives at,

$$r_h^4 = \left[\frac{\sqrt{(R^3 L_{iso}\rho_*)}\ \sigma_{p\gamma\to\Delta}\ \delta t\ \epsilon_p}{\sqrt{\pi}\times\left(.2\ \mathrm{GeV}^2\right)}\right]^{\frac{2}{5}}. \tag{3.7}$$

If we assume, the variability time $\delta t \sim 10^{-4}$ sec, $L_{iso} = 10^{51} L_{52}$ erg/sec, and the energy of the proton $\epsilon_p = 10^5$ GeV. Then the radial distance from the center to have pion formation, r_π turns out to be, $r_\pi \sim 10^{10.5}$ cm for $R = 10^{12}$cm and $\rho_* = 10^{-7}$ gm/cc. [1]. For more energetic protons, the radius of pion formation would shift further out-wards. For instance production of neutrinos of energy $\epsilon_\nu \geq 10^{17}$eV, would form further out. Once we have the estimate of the pion formation radius, we can estimate the strength of the magnetic field and find out the field induced modification to the spectra. The estimation of the magnetic field is done in the next subsection.

3.3 Magnetic Field Estimation

In this note we have assumed the system to follow ideal MHD conditions. That is to say that the magnetic energy density can be expressed following eqn. (2.4)

$$\left[\frac{B_o^2}{2}\right] \sim 2.1\cdot10^{53} L_{52}^{3/4}. \tag{3.8}$$

Where we have used the energy density at the injection point i.e., $r = r_o = 10^7$ cm with magnetic equipartition taken to be of order unity. We note that r_o is also the width of the shell, that the GRB wind expands with. Conservation of magnetic field energy at the wind rest frame implies at any radial distance r, in the GRB environment,

---

[1] We have used the fact $L_{iso}\delta t = 10^{51} L_{52}\mathrm{erg} = 0.6\times10^{55}\mathrm{L}_{52}$ GeV

$$4\pi r_o^2 B_o^2 = 4\pi r^2 B^2. \tag{3.9}$$

leading to the relation, $B \times r = B_o \times r_o$ implying $B = B_o\,(r_o/r)$. Using these relations, the magnetic field strength, in terms of GRB luminosity turns out to be, $B_o = \left[4.2.10^{53}L_{52}^{3/4}]^{1/2}\right]$. Hence the magnetic field strength at distance r from the center turns out to be,

$$B = \frac{r_o}{r}\left[4.2.10^{53}L_{52}^{3/4}\right]^{1/2}. \tag{3.10}$$

For conditions prevailing at the pion formation radius, corresponding to 10^5 GeV protons, the strength of the field– compared with the critical Field strength $B_c(simm_e^2)$– i.e $\frac{B}{B_{cr}} \sim 0.01$. The numbers mentioned here can get an order of magnitude variation depending on the details. Since the purpose of this note is to get an order of magnitude estimate of the effects, we reserve the details for future communications.

3.4 Estimates Of Energy Loss.

The high energy $\pi^+ \& \mu^+$ from the decay of Δ would undergo energy loss by inverse Compton (IC) scattering, if the optical depth is large. However even at low optical depth the end products remain coupled to photons provided the photon drag time $t_D \sim \frac{m_p}{\sigma_T u_\gamma}$ is comparable to their decay time [13]. One can verify that t_D is indeed comparable to the decay time, ensuring that the extremely relativistic pions and muons would undergo IC losses during their life time in the jet. However during the same period, they are expected to be accelerated in the reverse shocked plasma, to gain energy. Recalling the expression of pressure at the shock front to be $P_r = 4\left(\frac{\Gamma_j}{\Gamma_h}\right)^2 n_j m_p$, the equation of motion for the charged particles at the shock front would be given by:

$$\frac{dp}{dt} = 4\left(\frac{\Gamma_j}{2\Gamma_h}\right)^2 n_j m_j \times \ Area. \tag{3.11}$$

If the jet makes an angle θ, at a distance r from the center then the area would be $\pi(r\theta/2)^2$. One can further use the fact that $\frac{\theta}{\Gamma_j \sim 1}$, one gets,

$$\frac{dE_j}{E_j dt} = \frac{\pi n_j m_j r_o^2}{4E_j} = \frac{\pi n_j r_o^2}{4\Gamma_j}. \tag{3.12}$$

While coming to the relation above, we have made a conservative estimate, by replacing r by r_o and neglecting the rest mass of the particles. During the life time τ, the fraction of energy gained, by the particles would be,

$$\Delta E_j = \frac{\pi n_j r_o^2 E_j \tau}{4\Gamma_j}. \tag{3.13}$$

The inverse Compton energy loss, in the ultra relativistic regime has already been estimated in [12], and is given by

$$\Delta_{IC} = \frac{E_j \tau \sigma_T (m_e/m_j)^2 n_\gamma}{\gamma_j (E_\gamma/m_j)}. \tag{3.14}$$

Finally the ratio,$\frac{\Delta_{IC}}{\Delta_j} = \frac{4}{\pi}\frac{\sigma_T m_e^2 n_\gamma}{E_\gamma m_j n_j r_o^2}$. This ratio probably would decide whether the IC loss is significant enough to cause any significant suppression in neutrino production. For the kind of situation we are interested in, it seems this ratio would be less than unity hence, not much of suppression in the high energy neutrino spectrum. However this is a very tentative statement and one still needs to perform a quantitative estimation.

4 Discussion and Outlook.

High energy neutrinos moving in a magnetic field can decay to $\nu \rightarrow \nu l^+ l^-$. Where l can be either electron or muon or tau. The important point here is to note that, such processes are otherwise forbidden in vacuum. It has also been noted that, these processes are insensitive to neutrino mass and mixing in the leptonic sector. In a purely magnetic field the probablity of these processes depends on the parameter, $\mathcal{P} = \frac{eBp_\perp}{m_l^3}$. Depending on whether $\mathcal{P}$ is greater than unity or less than unity, the probablity for this reaction varies significantly. In a magnetized vacuum the probablity for such processes, α (for $\mathcal{P} >> 1$), is given by

$$\alpha_l = \frac{G_F^2 \left(g_v^2 + g_A^2\right) m_l^6 \mathcal{P}^2}{27\pi^2 E_\nu} \left[ln\mathcal{P} - \frac{1}{2} ln3 - \gamma_E - \frac{29}{21} - \frac{1}{\mathcal{P}^{\frac{2}{3}}} \frac{9}{56} \frac{3^{\frac{1}{3}}\pi^2}{\Gamma^4[\frac{2}{3}]} \frac{19g_V^2 - 63g_A^2}{g_V^2 + g_A^2} \right]. \tag{4.1}$$

We would like to remind here that γ_E is Euler's gamma function and $\Gamma[\frac{2}{3}]$ is the Gamma function. The last term in eqn. (4.1) contributes significantly when the neutrino flavor index and lepton flavor indices are different. we however would be interested in such a case.

In a GRB, the location of the neutrino production of a particular flavor strictly speaking would be different. However since the muon life time is pretty much small, therefore the location of ν production from π^+ decay and the same from μ^+ decay though would be different however the difference in length may not create significant changes in the medium properties. Therefore, for all practical purposes, we would assume them to originate from the same location. If the number density of ν of certain energy E_ν at the production point is given by $n_{0\nu_i}(E)$ then after travelling a distance dx the modified number density would be given by $dn_{\nu_i}(E) = -n_{0\nu_i}(E)\alpha dx$. And the solution to the same eqn., gives $n_{\nu_i(E)} = n_{0\nu_i}(E)e^{-\alpha_i x}$; where α_i is the flavor dependent absorption coefficient. We can further quantify the flavor dependent absorption as,

$$\kappa_i = -ln\left(\frac{n_{\nu_i(E)}}{n_{0\nu_i}(E)}\right) = \alpha_i x. \tag{4.2}$$

Therefore one can define the asymmetry parameter A_p, to be:

$$A_p = \frac{\kappa_{\nu_\mu} - \kappa_{\nu_e}}{\kappa_{\nu_\mu} + \kappa_{\nu_e}} = \frac{\alpha_{\nu_\mu} - \alpha_{\nu_e}}{\alpha_{\nu_\mu} + \alpha_{\nu_e}}. \tag{4.3}$$

A measure of this quantity A_p can in principle give us information about the nature of the medium at the production site of the neutrinos as well as the neutrino properties. In section [3.2] we had estimated the pion formation radius for $\epsilon_p \geq 10^5 GeV$ followed by the estimation of magnetic field at that location. It should be noted that any enhancement of the magnetic fields due to charge particle streaming has been ignored here. We assumed the magnetic field to be toroidal and the neutrino propagation direction to be perpendicular to it. Furthermore we assumed the strength of the magnetic field to be held constant over the thickness of the shell (in its rest frame) i,e $r_o = 10^7$cm. This can be justified because any variation to magnetic field over a distance larger than $\frac{1}{m_e}$ would not make any significant contribution to the probablity amplitude we are interested in.

Coming back to the question of asymmetry, as has already been discussed the asymmetry parameter A_p as has already been discussed, is far from zero. For very high energy neutrinos it seems to decrease compared to the low energy neutrinos, see fig.[4]. However estimation of actual degradation of energy spectrum of TeV or Super TeV Neutrinos would require more careful estimation of the parameters discussed here as well as taking matter effects into consideration. However there is an interesting consequence of this effect, that is, ther might be some enhancement in the number of neutrinos coming from the decay of muons originating from the reaction $\nu_e \rightarrow \nu_e \mu^+ \mu^-$. That might have interesting consequences for the upcoming neutrino experiments.



Figure 1: Asymmetry parameter (A_p, see text) for ν_e and ν_μ for the High Energy neutrinos in the GRB enviornment. x - Axis corresponds to neutrino energy in MeV and y Axis corresponds to A_p. Magnetic field strength in units of electron mass is taken to be 0.001. Energy is taken to be same for both the flavors.

5 acknowledgement

The author wishes to thank Profs. P.B. Pal and P. Bhattacharyya and K. Kar for discussions and exchange of views. He would also like to thank Dr. S. Razzaque for bringing few articles to his notice.

References

[1] Gulibert, P.W., Fabian, A,C., Rees, M.J., MNRS, **205** 593(1983). Belinnikov, S., astroph/9911138.

[2] M.J. Rees and P. Meszaros , Astrophys. J.430, L93 (1994).

[3] K. Bhattacharya, A. K. Ganguly and S. Konar, Phys. Rev. **D65** 013007,2002. K. Bhattacharya and Avijit K. Ganguly Phys. Rev. **D68** 053011 (2003). K. Bhattacharya and Avijit. K. Ganguly :hep-ph/0209237; and see the references there in. S.P. Mikheyev and A. Au.Smirnov, Sov.J. Nucl. Phys. For a detailed exposition to this topic one can see, K. Bhattacharyya and P. B. Pal. Proc. Indian. Natn Sci Acad. **70 A** No 1. January 2004, pp. 145-161.

42, 913 (1985). L. Wolfenstein, Phys. Rev **D17**, 2369 (1978).

[4] H. A. Bethe, Rev. Of. Mod. Phys. **62**, 1990, 801.

[5] J. N. Bahcall and P. Me'sza'ros, Phys. Rev. Lett. 85, 1362 (2000).

[6] E. Waxman and J. Bahcall, Phys. Rev. Lett. 78, 2292 (1997).

[7] P. Meszaros and E. Waxman and , Phys. Rev. Lett. 87, 171102-1 (2001).

[8] J. Tan, C. D. Matzner and C.F. McKee, Astrophys. **J** 946 (2001).

[9] Eli Waxman, astroph/0303517.

[10] M. J. Rees and P. Meszaros, Astrophys. J. Lett, 430, L93.

[11] see for instance reference [9]

[12] Soebur Razzaque, Peter Me'sza'ros and Eli Waxman, astroph/0303505

[13] P. Meszaros, P. Laguna and M. J. Rees, astro-ph/9301007.

[14] A. V . Kuznetsov, N. V. Mikheev, Phys. Lett. **B394** ,123 (197).